

How To Properly Apply The RAN™-Fentanyl Transdermal Patch.

Please read the RAN™-Fentanyl Transdermal System patch package insert for additional important safety information.



Step 1. PREPARING THE ADHESION SITE

RAN™-Fentanyl Transdermal patches should be applied to a clean, dry area of your chest, back, flank or upper arm. Clean the skin, using clear water only, and pat dry. Hair at the chosen application site should be clipped close to the skin (do not shave).



Step 5. WASH YOUR HANDS

When you have finished applying the patch wash your hands using water only. Do not use soap, alcohol, or other solvents to remove any gel residue because they may increase the drug's ability to go through the skin.



Step 2. REMOVE FROM THE POUCH

Each RAN™-Fentanyl Transdermal patch is sealed in a protective pouch. Do not remove the patch from the pouch until you are ready to use it. Tear open the pouch at the notched corner and remove the patch. Dispose of the pouch in your regular garbage.



Step 6. FILL OUT AND ATTACH THE DATE LABEL

Special date labels are provided to help you remember when to apply your next patch. After putting on the patch, write the current date and time on a label, then stick the label on the patch.



Step 3. REMOVE THE STICKY PAD COVER

A stiff protective liner covers the sticky side of the patch – the side that will be applied to your skin. Hold the liner at the edge and pull the patch from the liner. Try not to touch the sticky side of the patch. Discard the liner in your regular garbage.



Step 7. PATCH REMOVAL AND REPLACEMENT

After wearing the patch for 3 days, or as directed by your doctor, remove it and fold it in half so the sticky sides attach together. Flush the old, folded patch down the toilet immediately.

Choose a different location on your skin to apply a new patch and repeat steps 1 to 6 in order.



Step 4. APPLY TO SKIN SURFACE

Immediately after you have removed the liner, apply the sticky side of the patch to the prepared skin surface. Press the patch firmly on your skin with the palm of your hand and hold it there for about 30 seconds. If you have problems with the RAN™-Fentanyl Transdermal patch not sticking, apply first-aid tape to the edges of the patch only.

WHAT TO DO IF THE PATCH LOOSENS OR FALLS OFF?

If the patch falls off, dispose of the old one as directed above and apply a new one, making sure the new skin area you choose is clean, dry and free from hair.

Do not cut, damage or chew RAN™-Fentanyl Transdermal System patch. If it is damaged in any way it will not work properly, may not be safe, or may lead to possible uncontrolled release of fentanyl and life-threatening consequences.

RANBAXY

For more information call: 1-866-840-1340

TM – Trademark of Ranbaxy Laboratories Limited